

Adequacy of Current Observing Systems for Famine Early Warning

The breakout group began by reviewing the requirements and context of agricultural monitoring for famine early warning:

- EWS community has to outperform the media (CNN) in reaching the decision maker
- Need to use scientifically sound methods to present credible, convincing evidence that will evoke early response
- Need to focus on observations that constitute powerful, fundamental indicators

Adequacy of Current Observing Systems for Famine Early Warning

What observing systems are currently being used?

- Precipitation observations from stations, and satellite estimates
- Satellite vegetation index data

Are they adequate?

Precipitation

- Station observations are good, but data available through WMO GTS are too sparse
 - Station networks are more extensive than what is available via GTS
 - National met service policies prevent more complete release of data
 - Many food insecure regions, especially pastoral areas, lack coverage altogether
- Satellite estimates offer complete spatial coverage at high frequency (3hr - 24hr) in near real time, and are freely available over the web from several sources
 - Satellite estimates suffer from biases that can produce misleading results

Adequacy of Current Observing Systems for Famine Early Warning

What are prospects for continuity/enhancement?

Precipitation

- National met service 'cost-recovery' station data policies are unlikely to change any time soon
- Introduction of additional stations in food insecure regions faces obstacles of initial costs and maintenance, logistical and data management capacity
- Satellite estimates have good prospects for continuity/enhancement thanks to the international satellite constellation being implemented through the Global Precipitation Mission
- Satellite estimates can be enhanced through bias removal post-processing with (1) gridded station climatologies, and (2) blending of locally available, non-GTS station observations

Actions

- FEWS NET will identify 3-4 countries/regions where need is great and prospects are high for cooperation to create gridded climatologies and implement blending of locally available extra station data
- Approach WMO for endorsement of these activities, easing data access policies

Adequacy of Current Observing Systems for Famine Early Warning

Are they adequate?

Vegetation Indices

- NDVI from AVHRR has a long history of use in famine early warning, are produced regularly every 10-15 days, and are available freely in near real time over the web, and offer a long time series (since 1981)
- NDVI from AVHRR has seen decreasing utilization for famine early warning due to ambiguity of calculated greenness anomalies, loss of confidence
 - Removal of noise in data time series is difficult (sensor aging, orbital degradation), especially for dual gain instruments as on NOAA 17
- NDVI from MODIS has proven to be of excellent quality and well-suited to characterization of crop conditions, despite shorter time series (since 2000)
- Access to NDVI from MODIS through conventional DAAC procedures is problematic
 - Delays of several days following fixed compositing periods can mean data are 'old' for early warning purposes
 - Data tiles cannot be downloaded to some developing countries due to excessive file size versus available band width

Adequacy of Current Observing Systems for Famine Early Warning

What are prospects for continuity/enhancement?

Vegetation

- NASA/USDA/UMd are implementing a global, near real time system for MODIS NDVI
- No further MODIS missions are planned. Continuity with data from the VIIRS instrument subject to delays, with consequent risk of a data gap

Actions

- NASA/USDA/UMd investigate famine early warning requirements for spatial coverage and download file size
- Investigate alternative satellite platforms for follow-on instruments
- Investigate data from wide field of view instruments (India-AWiFS, CBERS, etc) as higher resolution sources of NDVI
- Investigate SAR imagery for detection of crop emergence